## Prevent Accidental Poisoning



Use this checklist to prevent accidental poisoning in your home.

## Medicines

- ☐ Store medicines out of children's reach and sight.
- ☐ Keep products in their original containers and read directions
- ☐ Don't take medicine or vitamins in front of children. Don't call them "candy."
- ☐ Keep visitors' purses, bags and coats out of the reach of children.
- ☐ Use only the dosing device that comes with the liquid medicine.
- □ When other caregivers are giving your child medicine, write clear instructions about what medicine to give, how much to give and when to give it.



## Cleaning Products Store cleaning products out of children's reach and sight. Keep products in their original containers and read product labels. ☐ Buy child-resistant packages when available. Never mix cleaning products. Carbon Monoxide Install carbon monoxide detectors in all П bedrooms and all levels of the home. Check the batteries in detectors every month. ☐ Check gas heaters, gas furnaces, fireplaces and wood-burning stoves for safety every fall. Lead Test for lead-based paint in homes built before 1978. Encourage children to play in grassy areas of the yard to decrease potential exposure to lead found in soil. Leave shoes outside the home to avoid spreading any lead dust from outside. Hire an expert if you need to remove lead. ☐ A balanced diet can help a child absorb less lead if exposed to it. If your child has been poisoned... Call 911 if the child is choking, can't breathe, or is having a seizure. Otherwise, call the poison control hotline, 1-800-222-1222, and be sure to have the product with you when you call.



Follow the operator's instructions. Instructions

are different depending on the item swallowed. Please don't make the child vomit or give them anything to drink unless

instructed.